

Original Article

Cite this article: Dixit S, Manjebayakath H, Saravanane N (2021). Two new *Pseudoceros* (Platyhelminthes: Polycladida: Pseudocerotidae) from Agatti Island, India and a species checklist from Indian waters. *Journal of the Marine Biological Association of the United Kingdom* 1–11. <https://doi.org/10.1017/S0025315421000151>

Received: 28 May 2020

Revised: 14 February 2021

Accepted: 22 February 2021

Key words:

Cotylea; Indian Ocean; marine flatworm; novel species; Rhabditophora; taxonomy

Author for correspondence:

Sudhanshu Dixit, E-mail: sid2130@gmail.com

Two new *Pseudoceros* (Platyhelminthes: Polycladida: Pseudocerotidae) from Agatti Island, India and a species checklist from Indian waters

Sudhanshu Dixit , Hashim Manjebayakath and Narayanane Saravanane

Centre for Marine Living Resources and Ecology, Atal Bhavan, Puthuvype, 682508, Kochi, India

Abstract

The Lakshadweep archipelago constitutes a major coral region of India but still lacks sufficient biodiversity data owing to its remoteness and a low number of faunal studies in the past. The present paper describes two new *Pseudoceros* species collected from Agatti Island, Lakshadweep, India. *Pseudoceros bipurpurea* sp. nov. and *Pseudoceros galaxea* sp. nov. are described based on external and internal characters, supported with histological studies and photographs. *Pseudoceros bipurpurea* sp. nov. is characterized by a cream background colour and an orange median line surrounded by dense patches of purple spots, which tend to disperse and broaden towards the posterior end. *Pseudoceros galaxea* sp. nov. displays a brown background colour with numerous white to cream dots covering almost the entire dorsal surface and a thin black margin. This study adds two new species to the polyclad fauna in Indian waters, raising the count to 68 species. An updated checklist of polyclads from Indian coast is also provided.

Introduction

The Lakshadweep archipelago consists of 36 islands, 10 of them inhabited, and five submerged reefs, the only coral atolls in India (Ghosh, 1991). The islands are scattered over a large stretch of the Arabian Sea, off the south-west coast of India (Figure 1). They are all characterized by the presence of a large lagoon with numerous reef patches on the western side, and steep slopes and rich coral reefs on the eastern side.

Polyclads of the suborder Cotylea, Lang, 1884 are characterized by the presence of a ventral sucker-like organ called a cotyl which is posterior to the female gonopore (Lang, 1884). Velasquez *et al.* (2018) indicated the presence of around 380 cotylean species worldwide, with highest diversity in the Caribbean and the Indo-Pacific regions. The genus *Pseudoceros* is the most populous genus under the suborder Cotylea with a total of 138 species worldwide (1 species, Jie *et al.*, 2016; 1 species, Cuadrado *et al.*, 2017; 135 species, Tyler *et al.*, 2006–2020; 1 species, Ramos-Sánchez *et al.*, 2020). Because of homogeneity in reproductive structures, some authors suggested that the specific identification relies on the colour and colouration pattern of the *Pseudoceros* species (Prudhoe, 1989; Newman & Cannon, 1994, 1996, 1997, 1998). This criterion of identification based on colours was further supported by Goggin & Newman (1996) and Litvaitis & Newman (2001) based on ribosomal nucleotide sequences. However, when species share relatively homogeneous external features such as colour or colouration patterns, internal characters such as orientation of seminal vesicle, morphology of the prostatic vesicle, depth of the male atrium, shape and size of the stylet, distance between sucker and female gonopore (Bolaños *et al.*, 2007; Marquina *et al.*, 2015) and musculature of the penile sheath (Ramos-Sánchez *et al.*, 2020) can be considered.

The first survey in Lakshadweep archipelago dated from the beginning of the 20th century (Ladilaw, 1902) and was carried out in Minicoy Island. The only available additional information about the polyclad fauna of these islands are reports from Kavaratti Island and Agatti Island (Apte & Pitale, 2011; Dixit *et al.*, 2019, 2021). This paper is a result of a continuous survey being undertaken at Agatti Island by CMLRE to assess the biodiversity of polyclads and describe two pseudocerotid flatworms. With this study, the number of polyclads reported from this archipelago will increase to 29 species. The present contribution not only adds new information to the biodiversity of polyclads in these unique ecosystems but also provides an updated checklist of polyclads inhabiting Indian waters (Table 1).

Materials and methods

Specimens were collected during scuba diving in subtidal and by hand picking in intertidal areas of Agatti Island (Figure 1) and photographed using an Olympus TG-5 camera with underwater housing. Animals were fixed on frozen 10% formalin buffered with seawater (modified methodology of Newman & Cannon, 2003). Key external morphological characters such as cerebral eyes, tentacular eyes, marginal eyes, pseudotentacles, dorsal and ventral surface were photographed with a stereo microscope (Leica, M80) and measurements such as total






Fig. 1. Lakshadweep Islands (black rectangle) near the west coast of southern India (top) and magnified image of Agatti Island (below) with collection sites of new species – *Pseudoceros galaxea* sp. nov. (red dot); *Pseudoceros bipurpurea* sp. nov. (yellow dot). Source: Google Earth.

length and width of the specimen, distance between mouth, gonopores and sucker were made. The portions with reproductive structures were dissected out, serially sectioned (7–8 μ m), stained with haematoxylin and eosin and photographed with a LEICA DMI8 inverted microscope. The holotypes and paratype of the newly described species are deposited at the Referral Centre at Centre for Marine Living Resources and Ecology (CMLRE), Kochi, India.

Results

SYSTEMATICS

Order POLYCLADIDA Lang, 1881
 Suborder COTYLEA Lang, 1884
 Family PSEUDOCEROTIDAE Lang, 1884
 Genus *Pseudoceros* Lang, 1884
Pseudoceros bipurpurea sp. nov. Dixit
 (Figures 2–4)

Type material

Holotype: One specimen, 16 $\times$ 9 mm, serial sectioned and mounted on 43 slides. Solar Point, Agatti Island, Lakshadweep, India (10°50'28"N 72°11'30"E); sandy substratum; water depth: 32 m; (CMLRE IO/DV/POY/00028); Coll. S. Dixit, 6 May 2019.

Etymology

The specific name *bipurpurea* is a compound Latin noun meaning 'two purple' refers to the characteristic dorsal colour pattern, two (*bi*) purple (*purpura* in Latin) regions of dense purple spots surrounding the orange median line.

Diagnosis

Background body colour cream with an orange median band surrounded by dense patches of purple spots without touching the margin (Figure 2A). These dense purple spots tend to disperse and broaden toward posterior end. Marginal rim blue.

Description

Live: Body oval and margin without ruffles. Background body colour cream with an orange median band. The median band is surrounded by dense purple spots (Figure 2A). These purple spots tend to disperse and broaden toward posterior end encircling the orange median band. The median band and purple spots start at the same point below cerebral eyes cluster and end without touching the posterior end. Marginal rim blue (Figure 2A). Pseudotentacles are simple folding of the anterior margin with same colour pattern as margin. Cerebral eyes dense and more or less forming a circular cluster (Figure 2A, B), tentacular eyes hard to locate due to the blue colour of the pseudotentacles and only visible in the preserved specimen (Figure 2B). The ventral surface is of cream colour.

Preserved: Specimen whitish and colour lost after fixation (Figure 2B, D). Orange median band discoloured and purple spots turned light brown showing the spotted pattern more clearly. Male and female gonopores are 1 mm apart, while female gonopore and sucker are 1.3 mm apart (Figure 2C). Pharynx ruffled with eight folds and male gonopore is situated very near to last pair of pharyngeal folds (Figure 2C). Pharynx is 3.2 mm long and distance between mouth and male gonopore is 2.3 mm. Cerebral eye cluster with about 60–70 eyes. Tentacular eyes arranged in two scattered lines (Figure 2B).

Reproductive system: The male copulatory apparatus consists of seminal vesicle, free prostatic vesicle, penis papilla armed with a stylet housed in male atrium which opens to the outside via male gonopore. An elongated seminal vesicle (670 μ m long and 280 μ m high) is present (Figures 3A & 4). An oval and thick-walled prostatic vesicle (192 long $\times$ 89 μ m high) is present anterior to seminal vesicle (Figures 3C & 4). Prostatic vesicle has smooth glandular lining, 12–15 μ m high (Figure 3C). Prostatic and ejaculatory duct enter separately in penis papilla and join before stylet (Figures 3D & 4). Male atrium conical (Figure 3C) housing conical penis papilla (130 μ m) armed with an elongated stylet (241 μ m) (Figure 3C, E) opening to the exterior via male gonopore (Figure 3C).

Female copulatory apparatus consists of oviduct, vagina, cement pouch, cement glands, female atrium and female gonopore (Figures 3B & 4). The common oviduct enters in vagina which then leads to a short female atrium via cement pouch. Numerous cement glands are scattered in parenchyma tissue from oviduct to seminal vesicle surrounding the vagina. Female atrium opens to exterior via female gonopore (Figures 3B & 4).

Taxonomic remarks

Presence of smooth dorsal surface, ruffled pharynx, cerebral and tentacular eyes, marginal tentacles formed by upfolding of anterior margin, centrally located sucker behind female gonopore, male copulatory apparatus single and just behind pharyngeal cavity, free prostatic vesicle and penis papilla armed with stylet place this new species within the family Pseudocerotidae and in the genus *Pseudoceros*.

Pseudoceros bipurpurea sp. nov. differs from all other *Pseudoceros* species by its colour pattern on dorsal surface. In terms of background colour and marginal rim (cream background and blue rim) the newly described species here matches with *Pseudoceros indicus* Newman & Schupp, 2002, *P. gamblei* Laidlaw, 1902, *P. rubrotentaculatus* Kaburaki, 1923 and *P. concinnus* (Collingwood, 1876) but differs in terms of colour pattern on dorsum. Several species of *Pseudoceros* display median




Fig. 2. *Pseudoceros bipurpurea* sp. nov. (A) Holotype (live); (B) cerebral and tentacular eyes (preserved specimen); (C) ventral surface (preserved specimen); (D) dorsal surface (preserved specimen). ce, cerebral eyes; fp, female gonopore; m, mouth; mp, male gonopore; p, pharynx; su, sucker; te, tentacular eyes.

stripes (either one or more than one). A species which possesses a single median stripe of matching colour (orange) on dorsum is *Pseudoceros galatheensis* Dixit, Raghunathan & Chandra, 2017 (thin orange median stripe) but it has light blue background colour which grows darker towards margins which differentiates it with newly described species here. However, there are a few species with median stripe bordered or surrounded by other stripes of different colour which include *Pseudoceros bifurcus* Prudhoe, 1989 (white median stripe outlined with dark burgundy), *Pseudoceros monostichos* Newman & Cannon, 1994 (black brown median line bordered by white and light brown), *Pseudoceros susanae* Newman & Anderson, 1997 (white median stripe surrounded by two orange stripes) and *Pseudoceros tristriatus* (three orange stripes on dorsum). *Pseudoceros bipurpurea* sp. nov. differs from all above-mentioned species by presence of an orange median stripe surrounded by dense purple spots. As no other valid species under the genus *Pseudoceros* is comparable with the newly described species in terms of colouration and pattern, we establish *Pseudoceros bipurpurea* sp. nov. as a new species to science.

Pseudoceros galaxea sp. nov. Dixit
(Figures 5–7).

Type material

Holotype: One specimen, 28 × 20 mm, serial sectioned and mounted on 22 slides. Intertidal area near CMLRE field research

station, Agatti Island, Lakshadweep, India (10°50'38"N 72°11'17"E) (CMLRE IO/IT/POY/00029); Coll. S. Dixit, 7 May 2019.

Paratype: One specimen, 16 × 12 mm. Location and collection details same as holotype (CMLRE IO/IT/POY/00030).

Etymology

The specific name *galaxea* is the Latin genitive form of *galaxiās* (galaxy), which means 'belong to the galaxy' and refers to the colour pattern resembling the stars in a galaxy.

Diagnosis

Background body colour chocolate brown with numerous small white to cream dots on the entire dorsal surface (Figure 5A). These minute dots are very densely distributed giving a spray-like appearance. Some white dots are too close forming clusters appearing like bigger dots. A black marginal band studded with white microdots runs around the whole body including pseudotentacles.

Description

Live: Body large, oval and margin slightly ruffled. Background body colour chocolate brown with numerous small white to cream dots on entire dorsal surface (Figure 5A). Median area slightly darker in colour and raised. White dots are densely distributed giving a spray-like appearance. Some white dots are




Fig. 3. *Pseudoceros bipurpurea* sp. nov. Sagittal sections (A) showing arrangement of reproductive structures; (B) female reproductive system; (C) and (D) male reproductive system; (E) stylet (magnified). cg, cement glands; cp, cement pouch; ed, ejaculatory duct; fa, female atrium; fp, female gonopore; ma, male atrium; mp, male gonopore; ov, oviduct; pd, prostatic duct; pp, penis papilla; ps, penis sheath; pv, prostatic vesicle; st, stylet; sv, seminal vesicle; va, vagina.




Fig. 4. *Pseudoceros bipurpurea* sp. nov. Diagrammatic reconstruction of male and female reproductive system. cg, cement glands; cp, cement pouch; ed, ejaculatory duct; fa, female atrium; fp, female gonopore; in, intestine; ma, male atrium; mp, male gonopore; ov, oviduct; p, pharynx; pd, prostatic duct; pp, penis papilla; ps, penis sheath; pv, prostatic vesicle; st, stylet; sv, seminal vesicle; va, vagina.

very close forming small white clusters delimiting bigger dots. Marginal rim black (Figure 5A, C) with white dots which are smaller than dots on dorsum. Pseudotentacles are simple folding of the anterior margin, black and spotted. Intensity of spots

decreases towards margin (Figure 5B). Cerebral eyes cluster horse-shoe shaped, tentacular eyes hard to recognize due to black colour of pseudotentacles however some eyes can be seen on ventral side of pseudotentacles. Ventral surface light brownish.




Fig. 5. *Pseudoceros galaxea* sp. nov. (A) Holotype (live); (B) pseudotentacles and cerebral eyes (preserved specimen); (C) ventral surface (preserved specimen). ce, cerebral eyes; fp, female gonopore; m, mouth; mp, male gonopore; p, pharynx; pt, pseudotentacles; su, sucker.

Preserved: Measurements based on holotype. Specimen brown in colour even after fixation but spots turned pale. Mouth and male gonopore are 2.9 mm, male and female gonopore are 2.2 mm while female gonopore and sucker are 2.8 mm apart (Figure 5C). Pharynx ruffled with nine folds (eight in paratype) and male gonopore is situated immediately behind pharynx. Cerebral eyes cluster with about 35–40 eyes (Figure 5B).

Reproductive system: The male copulatory apparatus consists of a seminal vesicle (Figure 6A), a free prostatic vesicle and an armed penis papilla (Figures 6 A, B & 7) housed in male atrium which opens outside via male gonopore. An oval seminal vesicle, $517 \times 257 \mu\text{m}$, its rounded part oriented towards prostatic vesicle while tapered part oriented towards cement glands (Figure 6A). A free, small and circular prostatic vesicle ($75 \times 48 \mu\text{m}$) present anterior to the seminal vesicle (Figure 6B). Thickness of prostatic vesicle's muscular wall varies from 10–14 μm . The male atrium is wide housing a reduced penis papilla armed with a stylet (197 μm) (Figure 6B). Female copulatory apparatus consists of oviduct, vagina, cement pouch surrounded by dense cement glands and female atrium (Figure 6C, D). The vagina is oriented towards posterior region of body connecting to oviducts. The vagina opens to a very short female atrium. Cement pouch wide which receives secretion from numerous cement glands distributed in parenchyma tissue. The female atrium opens to exterior via female gonopore (Figures 6C & 7).

Taxonomic remarks

The presence of characters such as ruffled pharynx, male copulatory apparatus right behind pharyngeal cavity, free prostatic vesicle, marginal tentacles formed by upfolding of anterior margin and centrally located sucker behind female gonopore (Faubel, 1984) matches with the characters of the family

Pseudocerotidae. The presence of characters such as smooth dorsal surface, single male copulatory apparatus with seminal vesicle and armed penis papilla, pseudotentacles as simple folds of anterior margin, female gonopore equidistant from male gonopore and sucker (Faubel, 1984; Newman & Cannon, 1998) matches the diagnosis of the genus *Pseudoceros*. The newly described species differs from all other congeners on the basis of spotted pattern and marginal rim. However, species such as *Pseudoceros astrorum* Bulnes & Torres, 2014, *Pseudoceros bicolor* Verrill, 1901, *Pseudoceros canadiensis* Hyman, 1953, *Pseudoceros josei* Newman & Cannon, 1994, *Pseudoceros kylie* Newman & Cannon, 1998 and *Pseudoceros stellans* Dixit, 2019 exhibit similar background colour, dorsally with blotches or spots of different sizes and marginal band. These species are compared with the newly described species below.

Pseudoceros astrorum, described from Brazil, is characterized by dark brown ground colour, net-like pattern of small black granules, white spots of different sizes uniformly distributed, a thin black submarginal band with white rim and anterior margin and cerebral region devoid of pigment. *Pseudoceros galaxea* sp. nov. differs from *P. astrorum* because of absence of the white rim, net-like pattern of black granules and pigment devoid area in cerebral region.

Pseudoceros bicolor, described from Bermuda, its dorsal colouration varies from yellow to dark brown with white spots on dorsum, white marginal band with greyish black transverse stripes. The marginal band in *P. galaxea* sp. nov. is black with no transverse stripes which differentiates it from the former species.

Pseudoceros canadiensis, described from Canada, is apparently brown or flecked with brown as per its original description. This species differs from *P. galaxea* sp. nov. which possesses white to cream dots on dorsum and black rim.




Fig. 6. *Pseudoceros galaxea* sp. nov. Sagittal sections (A) male reproductive structures, cement glands and pharynx; (B) male reproductive system (magnified); (C) female reproductive system showing cement glands, cement pouch and female atrium; (D) female reproductive structures showing vagina and oviduct. cg, cement glands; cp, cement pouch; ed, ejaculatory duct; fa, female atrium; fp, female gonopore; in, intestine; ma, male atrium; ov, oviduct; p, pharynx; ps, penis sheath; pv, prostatic vesicle; st, stylet; sv, seminal vesicle.

Pseudoceros josei, described from Papua New Guinea, is transparent black with numerous small and densely packed yellow spots which turn smaller and white towards the margin while in *P. galaxea* sp. nov., the spots are white to cream and a black marginal band is present.

Pseudoceros kylie, described from Australia, is dark brown in background with cream microdots but has a bright orange broken band just before the rim while the marginal band is black and continuous in *P. galaxea* sp. nov.

Pseudoceros stellans, described from India, has a thick black marginal band with white to yellow spots and yellow blotches; median area marbled with irregular white shading while *P. galaxea* sp. nov. has a thin black marginal rim and dorsum is without any yellow blotches or shading.

As no other species was found comparable in the genus, in light of the above-mentioned characters and comparisons, *P. galaxea* sp. nov. is described as a new species to science.

Discussion

All the studies on the polyclads living in Indian waters have been conducted in the last nine years, except one study conducted a century ago by Laidlaw (1902). A total of nine species were

reported from Laidlaw's work which marked the beginning of polyclad research in India. A few species such as *Acanthozoon plenhi*, *Latocestus maldivensis*, *Planocera langi*, *Pseudoceros buskii* and *Pseudoceros tigrinus* described by Laidlaw from Lakshadweep have never been recorded since. These species are in need of re-description based on fresh samples and coloured photographs of internal and external details. From the year 1902 until 2011, not even a single study was conducted on these worms from Indian waters, including in the Lakshadweep Islands which are one of the major coral reef areas of India and nurture rich reef-associated biodiversity. However, many works were published on polyclads during this period from nearby geographic areas in Arabian Sea such as Sri Lanka (Kelaart, 1858; Laidlaw, 1904), Maldives (Laidlaw, 1902, 1903), Djibouti (Mexiner, 1907a, 1907b) and the Red Sea (Meyer, 1922; Prudhoe, 1952). Since 2011, efforts from a handful of researchers raised the number of species from nine to 68 from Indian waters (Apte & Pitale, 2011; Sreeraj & Raghunathan, 2011, 2013, 2015; Dixit & Raghunathan, 2013; Pitale *et al.*, 2014; Sreeraj *et al.*, 2015; Dixit *et al.*, 2015, 2017a, 2017b, 2018a, 2018b, 2018c, 2019, 2021; Pitale & Apte, 2017, 2019; Dixit, 2018; present study). All species of polyclads reported from Indian waters are compiled in a checklist (Table 1) based on published reports. The maximum number

Table 1. Checklist of polyclads reported from Indian coasts

	Species name	Region	References
	Order: Polycladida Lang, 1881		
	Suborder Cotylea Lang, 1884		
	Family: Pseudocerotidae Lang, 1884		
	Genus: <i>Acanthozoon</i> Collingwood, 1876		
1	<i>Acanthozoon alderi</i> (Collingwood, 1876)	AN	Dixit <i>et al.</i> (2018c)
2	<i>Acanthozoon fuscobulbosum</i> Dixit, Sivaperuman & Raghunathan, 2018	AN	Dixit <i>et al.</i> (2018a)
3	<i>Acanthozoon plehni</i> (Laidlaw, 1902)	LK	Laidlaw (1902)
	Genus: <i>Bulaceros</i> Newman & Cannon, 1996		
4	<i>Bulaceros newcannorum</i> Dixit, 2021	LK	Dixit <i>et al.</i> (2021)
5	<i>Bulaceros porcellanus</i> Newman & Cannon, 1996	LK	Dixit <i>et al.</i> (2021)
	Genus: <i>Phrikoceros</i> Newman & Cannon, 1996		
6	<i>Phrikoceros fritillus</i> Newman & Cannon, 1996	AN	Sreeraj <i>et al.</i> (2015)
7	<i>Phrikoceros katoi</i> Newman & Cannon, 1996	AN	Sreeraj <i>et al.</i> (2015)
8	<i>Phrikoceros mopsus</i> (Marcus, 1952)	AN	Sreeraj & Raghunathan (2015)
9	<i>Phrikoceros lizardensis</i> (Newman & Cannon, 1996)	AN, GOM	Dixit <i>et al.</i> (2015); Dixit (2018); Shrinivaasu <i>et al.</i> (2018)
	Genus: <i>Pseudobiceros</i> Faubel, 1984		
10	<i>Pseudobiceros apricus</i> Newman & Cannon, 1994	AN, LK	Sreeraj <i>et al.</i> (2015); Dixit <i>et al.</i> (2019)
11	<i>Pseudobiceros bedfordi</i> (Laidlaw, 1903)	AN	Sreeraj & Raghunathan (2013); Dixit (2018)
12	<i>Pseudobiceros damawan</i> Newman & Cannon, 1994	AN	Sreeraj & Raghunathan (2011); Dixit (2018)
13	<i>Pseudobiceros flavocanthus</i> Newman & Cannon, 1994	LK	Sreeraj & Raghunathan (2011)
14	<i>Pseudobiceros fulgor</i> Newman & Cannon, 1994	AN	Sreeraj <i>et al.</i> (2015); Dixit (2018)
15	<i>Pseudobiceros gratus</i> (Kato, 1937)	LK, AN	Apte & Pitale (2011); Dixit (2018)
16	<i>Pseudobiceros hancockanus</i> (Collingwood, 1876)	LK, AN, GOM	Apte & Pitale (2011); Dixit (2018); Shrinivaasu <i>et al.</i> (2018)
17	<i>Pseudobiceros kryptos</i> Newman & Cannon, 1997	AN	Dixit (2018)
18	<i>Pseudobiceros murinus</i> Newman & Cannon, 1997	LK, AN	Apte & Pitale (2011); Dixit (2018)
19	<i>Pseudobiceros splendidus</i> (Lang, 1884)	AN, LK	Sreeraj & Raghunathan (2013); Dixit (2018); Dixit <i>et al.</i> (2019)
20	<i>Pseudobiceros stellae</i> Newman & Cannon, 1994	LK, AN	Apte & Pitale (2011); Sreeraj & Raghunathan (2015)
	Genus: <i>Pseudoceros</i> Lang, 1884		
21	<i>Pseudoceros agattiensis</i> Dixit, 2019	LK	Dixit <i>et al.</i> (2019)
22	<i>Pseudoceros auranticrinis</i> Dixit, Raghunathan & Chandra, 2017	AN	Dixit <i>et al.</i> (2017b); Dixit (2018)
23	<i>Pseudoceros bicolor</i> Verill, 1901	LK	Dixit <i>et al.</i> (2019)
24	<i>Pseudoceros bifurcus</i> Prudhoe, 1989	AN	Sreeraj & Raghunathan (2011); Dixit (2018)
25	<i>Pseudoceros bipurpurea</i> sp. nov.	LK	Present study
26	<i>Pseudoceros bolool</i> Newman & Cannon, 1994	AN, LK	Sreeraj <i>et al.</i> (2015); Dixit (2018); Dixit <i>et al.</i> (2019)
27	<i>Pseudoceros buskii</i> (Collingwood, 1876)	LK	Laidlaw (1902)
28	<i>Pseudoceros concinnus</i> (Collingwood, 1876)	AN	Sreeraj & Raghunathan (2011); Dixit (2018)
29	<i>Pseudoceros cruentus</i> Newman & Cannon, 1998	AN	Dixit <i>et al.</i> (2015); Dixit (2018)
30	<i>Pseudoceros duplicinctus</i> Prudhoe, 1989	LK, AN, GOM	Apte & Pitale (2011); Dixit (2018); Shrinivaasu <i>et al.</i> (2018)
31	<i>Pseudoceros flavomarginatus</i> Laidlaw, 1902	LK	Laidlaw (1902)
32	<i>Pseudoceros galattheensis</i> Dixit, Raghunathan & Chandra, 2017	AN	Dixit <i>et al.</i> (2017a); Dixit (2018)
33	<i>Pseudoceros galaxea</i> sp. nov.	LK	Present study
34	<i>Pseudoceros gamblei</i> Laidlaw, 1902	AN	Sreeraj & Raghunathan (2011); Dixit (2018)
35	<i>Pseudoceros goslineri</i> Newman & Cannon, 1994	LK, AN	Apte & Pitale (2011); Sreeraj & Raghunathan (2011); Dixit (2018)

(Continued)

Table 1. (Continued.)

	Species name	Region	References
36	<i>Pseudoceros imitatus</i> Newman, Cannon & Brunckhorst, 1994	AN	Sreeraj <i>et al.</i> (2015); Dixit (2018)
37	<i>Pseudoceros indicus</i> Newman & Schupp, 2002	LK, AN, GOM	Apte & Pitale (2011); Sreeraj & Raghunathan (2013); Shrinivaasu <i>et al.</i> (2018)
38	<i>Pseudoceros intermittus</i> Newman & Cannon, 1995	AN	Sreeraj & Raghunathan (2013)
39	<i>Pseudoceros irretitus</i> Newman & Cannon, 1998	AN	Dixit <i>et al.</i> (2015); Dixit (2018)
40	<i>Pseudoceros jebborum</i> Newman & Cannon, 1994	AN	Dixit (2018)
41	<i>Pseudoceros leptostictus</i> Bock, 1913	AN	Sreeraj <i>et al.</i> (2015); Dixit (2018)
42	<i>Pseudoceros meenae</i> Dixit, Sivaperuman & Raghunathan, 2018	AN	Dixit <i>et al.</i> (2018a); Dixit (2018)
43	<i>Pseudoceros nigropunctatus</i> Dixit, Raghunathan & Chandra, 2017	AN	Dixit <i>et al.</i> (2017a); Dixit (2018)
44	<i>Pseudoceros paralaticlavus</i> Newman & Cannon, 1994	LK, AN	Apte & Pitale (2011); Dixit (2018)
45	<i>Pseudoceros rubronanus</i> Newman & Cannon, 1998	AN	Sreeraj & Raghunathan (2013)
46	<i>Pseudoceros scintillatus</i> Newman & Cannon, 1994	AN	Sreeraj <i>et al.</i> (2015); Dixit (2018)
47	<i>Pseudoceros</i> cf. <i>susanae</i> Newman & Anderson, 1997	LK	Apte & Pitale (2011)
48	<i>Pseudoceros stellans</i> Dixit, 2019	LK	Dixit <i>et al.</i> (2019)
49	<i>Pseudoceros stimpsoni</i> Newman & Cannon, 1998	AN	Sreeraj <i>et al.</i> (2015); Dixit (2018)
50	<i>Pseudoceros tigrinus</i> Laidlaw, 1902	LK	Laidlaw (1902)
51	<i>Pseudoceros tristratus</i> Hyman, 1959	AN	Sreeraj & Raghunathan (2013); Dixit (2018)
52	<i>Pseudoceros vishnui</i> Dixit, Raghunathan & Chandra, 2017	AN	Dixit <i>et al.</i> (2017b); Dixit (2018)
	Genus: <i>Thysanozoon</i> Collingwood, 1876		
53	<i>Thysanozoon brocchii</i> Risso, 1818	MH	Pitale & Apte (2017)
54	<i>Thysanozoon nigropapillosum</i> (Hyman, 1959)	AN	Sreeraj & Raghunathan (2013); Dixit (2018)
	Family: Euryleptidae Lang, 1884		
	Genus: <i>Cycloporus</i> Lang, 1884		
55	<i>Cycloporus australis</i> Prudhoe, 1982	MH	Pitale & Apte (2019)
56	<i>Cycloporus reticulatus</i> Newman & Cannon, 2002	MH	Pitale & Apte (2019)
57	<i>Cycloporus variegatus</i> Kato, 1934	MH	Pitale & Apte (2019)
58	<i>Cycloporus venetus</i> Newman & Cannon, 2002	AN	Sreeraj & Raghunathan (2015)
	Genus: <i>Eurylepta</i> Ehrenberg, 1831		
59	<i>Eurylepta aurantiaca</i> Heath & Mc Gregor, 1912–1913	MH	Pitale & Apte (2019)
	Genus: <i>Maritigrella</i> Newman & Cannon, 2000		
60	<i>Maritigrella fuscopunctata</i> Newman & Cannon, 2000	LK	Apte & Pitale (2011)
	Genus: <i>Prostheceraeus</i> Schmarda, 1959		
61	<i>Prostheceraeus fuscolineatus</i> Dixit, Raghunathan & Chandra, 2017	AN	Dixit <i>et al.</i> (2017b)
	Genus: <i>Stylostomum</i> Lang, 1884		
62	<i>Stylostomum mixtomaculatum</i> Pitale & Apte, 2019	MH	Pitale & Apte (2019)
	Suborder: Acotylea Lang, 1884		
	Family: Planoceridae Lang, 1844		
	Genus: <i>Paraplanocera</i> Ladilaw, 1903		
63	<i>Paraplanocera langi</i> (Ladilaw, 1902)	LK	Laidlaw (1902)
64	<i>Paraplanocera oligoglana</i> Schmarda, 1859	AN	Sreeraj <i>et al.</i> (2015)
	Family: Prosthiostomidae Lang, 1884		
	Genus: <i>Prosthiostomum</i> De Quaterfage, 1845		
65	<i>Prosthiostomum trilineatum</i> Yeri & Kaburaki, 1920	MH, AN	Pitale <i>et al.</i> (2014); Sreeraj <i>et al.</i> (2015)
	Family: Leptoplanidae Stimpson, 1857		

(Continued)

Table 1. (Continued.)

	Species name	Region	References
	Genus: <i>Euplanoida</i> Faubel, 1983		
66	<i>Euplanoida pardalis</i> (Laidlaw, 1902)	LK	Laidlaw (1902)
	Family: Diposthidae Woodworth, 1898		
	Genus: <i>Pericelis</i> Laidlaw, 1902		
67	<i>Pericelis byerleyana</i> (Collingwood, 1876)	LK	Laidlaw (1902)
	Family: Latocestidae Laidlaw, 1903		
	Genus: <i>Latocestus</i> Plehn, 1896		
68	<i>Latocestus maldivensis</i> (Laidlaw, 1902)	LK	Laidlaw (1902)

AN, Andaman and Nicobar Islands; GOM, Gulf of Mannar; LK, Lakshadweep Islands; MH, Maharashtra.




Fig. 7. *Pseudoceros galaxea* sp. nov. Diagrammatic reconstruction of male and female reproductive system. cg, cement glands; cp, cement pouch; ed, ejaculatory duct; fa, female atrium; fp, female gonopore; in, intestine; ma, male atrium; ov, oviduct; pd, prostatic duct; pp, penis papilla; ps, penis sheath; pv, prostatic vesicle; st, stylet; sv, seminal vesicle; va, vagina.

of species is reported from Andaman and Nicobar Islands (44 species) followed by Lakshadweep Islands (29 species). The genus *Pseudoceros* is dominating with 32 species in Indian waters out of which 14 species are from Lakshadweep waters. The genus *Pseudobiceros* follows next with 11 species from India while seven species are reported from Lakshadweep Islands alone. Other genera such as *Euplanoida* Faubel, 1983; *Eurylepta* Ehrenberg, 1831; *Latocestus* Plehn, 1896; *Maritigrella* Newman & Cannon, 2000; *Pericelis* Laidlaw, 1902; *Prosthoceraeus* Schmarda, 1859; *Prothiostomum* De Quaterfage, 1845; and *Stylostomum* Lang, 1884 are represented by only one species so far from Indian waters. Apte & Pitale (2011) reported *Pseudoceros* cf. *susanae* Newman & Anderson, 1997 from Lakshadweep Islands. This species though doubtful in its identification is retained in the checklist until new specimens of this species are collected for confirmation of its identification.

The discovery of five new species (two species from Dixit *et al.*, 2019; one species from Dixit *et al.*, 2021 and two from the present study) from this archipelago in recent times is a clear indication that more efforts are required for inventorization of polyclad fauna from these islands. Their extreme fragility has made these animals difficult to study (Newman & Cannon, 2003). During our field surveys, many polyclad specimens of different genera were collected, but some of them tended to curl, distort or dissolve during the course of fixation (personal observation). However, members of the genus *Pseudoceros* were less difficult to fix in comparison with species from other genera such as *Acanthozoon*, *Bulaceros*, *Thysanozoon* and *Pericelis* (personal observation). Thus, based on our observations, we speculate this as one of the reasons for a higher number of species being described under the genus *Pseudoceros* in comparison with other genera under suborder Cotylea. The flamboyant colouration

of species in the genus *Pseudoceros* may also be the reason for more numerous descriptions but more studies are required to test these speculations. We hope that recent publications in polyclad taxonomy from Indian waters will attract the interest of young researchers towards these extraordinary worms. The main food source of polyclads are oysters, bivalves and ascidians (Newman & Cannon, 2003), which are present in both intertidal and subtidal zones making these islands an ideal habitats for polyclads. Only three islands of the Lakshadweep archipelago have been surveyed for polyclad fauna to date (Agatti, Kavaratti and Minicoy), leaving 33 islands unexplored.

Acknowledgements. The authors would like to thank the Ministry of Earth Sciences (MoES), Government of India for providing the opportunity and facilities to visit Agatti Island. This work would not have been completed without support of the local staff of CMLRE field station at Agatti Island. The first author would like to thank Mr B. Kishore Kumar (CMLRE) for his support in logistics and collection of specimens. The first author would also like to thank his fellow colleagues from CMLRE for their help during fieldwork and comments during manuscript preparation. The support of three reviewers with their critical comments and in improving the manuscript is also acknowledged. This is CMLRE contribution number 132.

References

- Apte D and Pitale RD (2011) New records of polyclad flatworms (Platyhelminthes: Turbellaria) from coral reefs of Lakshadweep Island, India. *Journal of the Bombay Natural History Society* **108**, 109–113.
- Bock S (1913) Studien über Polycladen. *Zoologiska Bijdrag fran Uppsala* **2**, 31–344.
- Bolanos DM, Quiroga SY and Litvaitis MK (2007) Five new species of cotylean flatworms (Platyhelminthes: Polycladida) from the wider Caribbean. *Zootaxa* **1650**, 1–23.
- Bulnes VN and Torres Y (2014) *Pseudoceros astrorum*, a new species of Polycladida (Cotylea, Pseudocerotidae) from Northeastern Brazil. *Zootaxa* **3881**, 94–100.
- Collingwood C (1876) On thirty-one species of marine planarians, collected partly by the late Dr Kelaart, F.L.S. at Trincomalee, and partly by Dr Collingwood, F.L.S. in the eastern seas. *Transactions of the Linnean Society of London, Series 2, Zoology* **1**, 83–98.
- Cuadrado D, Moro L and Noreña C (2017) The Polycladida (Platyhelminthes) of the Canary Islands. New genus, species and records. *Zootaxa* **4312**, 38–68.
- De Quatrefages A (1845) Études sur les types inférieurs de l'embranchement des annelés: mémoire sur quelques planaires marines appartenant aux genres *Tricelis* (Ehr.), *Polycelis* (Ehr.), *Prosthiostomum* (Nob.), *Proceros* (Nob.), *Eolidiceros* (Nob.), et *Stylochus* (Ehr.). *Annales des Sciences Naturelles* **4**, 129–184.
- Dixit S (2018) *Taxonomy and Distribution of Family Pseudocerotidae (Platyhelminthes: Polycladida), Andaman and Nicobar Islands, India*. PhD thesis, Pondicherry University, India.
- Dixit S and Raghunathan C (2013) Polyclads of Andaman and Nicobar Islands. *Journal of the Andaman Science Association* **18**, 165–169.
- Dixit S, Sivaperuman C and Raghunathan C (2015) Three new records of polyclad flatworms from India. *Marine Biodiversity Records* **8**, e29.
- Dixit S, Raghunathan C and Chandra K (2017a) Two new marine flatworms (Polycladida: Pseudocerotidae) from Andaman & Nicobar Islands, India. *Zootaxa* **4221**, 111–122.
- Dixit S, Raghunathan C and Chandra K (2017b) Two new *Pseudoceros* (Polycladida: Pseudocerotidae) and a *Prostheceraeus* (Polycladida: Euryleptidae) from Andaman and Nicobar Islands, India. *Zootaxa* **4269**, 495–512.
- Dixit S, Sivaperuman C and Raghunathan C (2018a) Description of two new pseudocerotids (Rhabditophora: Rhabditophora; Polycladida) from Andaman & Nicobar Islands, India. *Zootaxa* **4403**, 365–377.
- Dixit S, Sivaperuman C and Raghunathan C (2018b) Erratum. *Zootaxa* **4442**, 600.
- Dixit S, Bulnes VN and Raghunathan C (2018c) Neotype designation for the marine flatworm, *Acanthozoon alderi* (Polycladida: Cotylea: Pseudocerotidae), from India with comments on the taxonomical status of the genus. *Zoological Studies* **57**, 45.
- Dixit S, Bayyana S, Manjebayakat H, Saravanane N and Sudhakar M (2019) Polyclad fauna of Agatti Island, Lakshadweep, India: new records and description of two new species. *Zootaxa* **4657**, 246–260.
- Dixit S, Manjebayakat H and Saravanane N (2021) A rare polyclad genus *Bulaceros* (Platyhelminthes: Polycladida: Pseudocerotidae): new species and new record from Indian coral atolls. *Marine Biology Research* **16**, 632–642.
- Ehrenberg CG (1831) Phytozoa, Turbellaria africana et asiatica. In Hemprich PC and Ehrenberg CG (eds), “*Symbolae physicae*.” *Animalia evertabrata exclusis insectis recensuit Dr. CG Ehrenberg. Series prima cum tabularum decade prima. Berolini, Fol. Phytozoa Turbellaria folia a–d, T4–5* [plates 4, 5 published in 1828].
- Faubel A (1983) The Polycladida, Turbellaria. Proposal and establishment of a new system. Part I. The Acotylea. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* **80**, 17–121.
- Faubel A (1984) The Polycladida, Turbellaria. Proposal and establishment of a new system. Part II. The Cotylea. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* **81**, 189–259.
- Ghosh AK (1991) Fauna of Lakshadweep: an overview. *Fauna of Lakshadweep: State Fauna Series 2. Zoological survey of India*. http://faunaofindia.nic.in/php/sfs/sfs_books_toc.php?book_id=005&type=sfs&book_title=Fauna+of+Lakshadweep.
- Goggin CL and Newman LJ (1996) Use of molecular data to discriminate pseudocerotid turbellarians. *Journal of Helminthology* **70**, 123–126.
- Heath H and McGregor E (1912–1913) New polyclads from Monterey Bay, California. *Proceedings of the Academy of Natural Science of Philadelphia* **64**, 455–488.
- Hyman LH (1953) The polyclad flatworms of the Pacific coast of North America. *Bulletin of the American Museum of Natural History* **100**, 265–392.
- Hyman LH (1959) A further study of Micronesian polyclad flatworms. *Proceedings of the United States National Museum* **108**, 543–597.
- Jie WB, Gan BQ, Cgen VY and Kuo SH (2016) *Pseudoceros magangensis*: a new species of pseudocerotid flatworm (Platyhelminthes: Polycladida) from Taiwan. *Platax* **13**, 33–50.
- Kaburaki T (1923) The polyclad turbellarians from the Philippine Islands. *Bulletin of the United States National Museum* **100**, 635–649.
- Kato K (1934) Polyclad turbellarians from the neighborhood of the Mitsui Institute of Marine Biology. *Japanese Journal of Zoology* **6**, 123–129.
- Kato K (1937) Polyclads collected in Idu, Japan. *Japanese Journal of Zoology* **7**, 211–231.
- Kelaart EF (1858) Description of new and little known species of Ceylon nudibranchiate molluscs and zoophytes. *Journal of the Ceylon Branch of the Royal Asiatic Society* **3**, 84–139.
- Laidlaw FF (1902) The marine Turbellaria with an account of the anatomy of some species. In Gardiner JS (ed.), *The Fauna and Geography of the Maldives and Laccadive Archipelagoes: Being the Account of the Work Carried on and of the Collections Made by an Expedition During the Years 1899 and 1900*, vol. 1. Cambridge: Cambridge University Press, pp. 282–312.
- Laidlaw FF (1903) On a collection of Turbellaria Polycladida from the Straits of Malacca. (Skeat Expedition 1899–1900). *Proceedings of the Zoological Society of London* **1**, 301–318.
- Laidlaw FF (1904) Report on the polyclad Turbellaria collected by Professor Herdman at Ceylon, in 1902. *Royal Society Report to Ceylon Government on Pearl Oyster Fisheries of Gulf of Mannar Part 2*, 127–136. <https://www.worldcat.org/title/report-on-the-polyclad-turbellaria-collected-by-professor-herdman-at-ceylon-in-1902/oclc/77785592>.
- Lang A (1881) Der Bau von Gunda segmentata und die Verwandtschaft der Plathelminthen mit. *Coelenteraten und Hirudineen* **3**, 187–251.
- Lang A (1884) Die Polycladen (Seeplanarien) des Golfes von Neapel und der angrenzenden Meeresabschnitte. Eine Monographie. *Fauna und Flora des Golfes von Neapel* **11**. Leipzig: Engelmann. <https://doi.org/10.5962/bhl.title.10545>.
- Litvaitis MK and Newman LJ (2001) A molecular framework for the phylogeny of Pseudocerotidae (Platyhelminthes, Polycladida). *Hydrobiologia* **444**, 177–182.
- Marcus E (1952) *Turbellaria brasileiros* (10). *Boletim da Faculdade de Filosofia, Ciências e Letras da Universidade de São Paulo. Zoologia* **17**, 5–186.
- Marquina D, Aguado MT and Noreña C (2015) New records of Cotylea (Polycladida, Platyhelminthes) from Lizard Island, Great Barrier Reef, Australia, with remarks on the distribution of the *Pseudoceros* Lang, 1884 and *Pseudobiceros* Faubel, 1984 species of the Indo-Pacific Marine Region. *Zootaxa* **4019**, 354–377.

- Meixner A** (1907a) Polykladen von der Somalikuest, nebst einer Revision der Stylochinen. *Zeitschrift für Wissenschaftliche Zoologie* **88**, 385–498.
- Meixner A** (1907b) Polyklades recueillis par M. Ch. Gravier dans le golfe de Tadjourah en 1904. *Bulletin du Muséum d'histoire naturelle Paris* **2**, 164–172.
- Meyer F** (1922) Polykladen von Koseir (Rotes Meer). (Kollektion Professor Klunzinger.) *Archiv für Naturgeschichte Abteilung A* **87**, 138–158.
- Newman LJ and Cannon LRG** (1994) *Pseudoceros* and *Pseudobiceros* (Platyhelminthes, Polycladida, Pseudocerotidae) from eastern Australia and Papua New Guinea. *Memoirs of the Queensland Museum* **37**, 205–266.
- Newman LJ and Cannon LRG** (1996) New genera of pseudocerotid flatworms (Platyhelminthes, Polycladida) from Australian and Papua New Guinean coral reefs. *Journal of Natural History* **30**, 1425–1441.
- Newman LJ and Anderson C** (1997) A new polyclad flatworm from the Maldives. *Journal of South Asian Natural History* **2**, 237–245.
- Newman LJ and Cannon LRG** (1997) Nine new species of *Pseudobiceros* (Platyhelminthes: Polycladida) from the Indo-Pacific. *Raffles Bulletin of Zoology* **45**, 341–368.
- Newman LJ and Cannon LRG** (1998) *Pseudoceros* (Platyhelminthes: Polycladida) from the Indo-Pacific with twelve new species from the Australia and Papua New Guinea. *Raffles Bulletin of Zoology* **46**, 293–323.
- Newman LJ and Cannon LRG** (2000) A new genus of euryleptid flatworm (Platyhelminthes, Polycladida) from the Indo-Pacific. *Journal of Natural History* **34**, 191–205.
- Newman LJ and Schupp P** (2002) A new pseudocerotid flatworm (Platyhelminthes, Polycladida) from the Indo-Pacific. *Micronesica* **34**, 177–184.
- Newman LJ and Cannon LRG** (2003) *Marine Flatworms. The World of Polyclads*. Clayton: CSIRO Publishing, 97 pp.
- Pitale R and Apte D** (2017) First record of *Thysanozoon brocchii* (Platyhelminthes: Polycladida) from Indian waters. *Marine Biodiversity Records* **10**, 1–11.
- Pitale R and Apte D** (2019) Intertidal euryleptid polyclads and description of a new *Stylostomum* Lang, 1884 from Maharashtra, India. *Zootaxa* **4652**, 317–339.
- Pitale R, Bhawe V and Apte D** (2014) First record of family Prosthlostomidae and *Prosthlostomum trilineatum* (Platyhelminthes: Polycladida) from the west coast of India. *Marine Biodiversity Records* **7**, 1–6.
- Plehn M** (1896) Neue Polykladen, gesammelt von Herrn Kapitän Chierchia bei der Erdumschiffung der Korvette Vettor Pisani, von Herrn Prof. Dr. Kükenthal im nördlichen Eismeer und von Herrn Prof. Dr. Semon in Java. *Jenaische Zeitschrift* **30**, 137–181.
- Prudhoe S** (1952) Manihine Expedition to the Gulf of Aqaba, 1948–1949. IV. Turbellaria: Polycladida. *British Museum (Natural History) Bulletin, Zoology* **1**, 175–179.
- Prudhoe S** (1982) Polyclad turbellarians from the southern coasts of Australia. *Records of the South Australian Museum (Adelaide)* **18**, 361–384.
- Prudhoe S** (1989) Polyclad turbellarians recorded from African waters. *Bulletin of the British Museum of Natural History, Zoology* **55**, 47–96.
- Ramos-Sanchez M, Bahia J and Bastida-Zavala R** (2020) Five new species of cotylean flatworms (Platyhelminthes: Polycladida: Cotylea) from Oaxaca, southern Mexican Pacific. *Zootaxa* **4819**, 49–83.
- Risso A** (1818) Mémoire sur quelques Gastéropodes nouveaux Nudibranches et Tectibranches observés dans la mer de Nice. *Journal de physique, de chimie, d'histoire naturelle et des arts* **87**, 272–376.
- Schmarda LK** (1859) Neue wirbellose Thiere beobachtet und gesammelt auf einer Reise um die Erde 1853 bis 1857. Bd I: *Neue Turbellarien, Rotatorien und Anneliden*. W. Engelmann, Leipzig. <https://doi.org/10.5962/bhl.title.85313>.
- Shrinivaasu S, Venkataraman K and Venkatraman C** (2018) Cryptofaunal pseudocerotid polyclads of Gulf of Mannar. *Global Journal of Bioscience and Biotechnology* **7**, 19–23.
- Sreeraj CR and Raghunathan C** (2011) New records of pseudocerotid polyclads from Andaman and Nicobar Islands, India. *Marine Biodiversity Records* **4**, e73.
- Sreeraj CR and Raghunathan C** (2013) Pseudocerotid polyclads (Platyhelminthes, Turbellaria, Polycladida) from Andaman and Nicobar Islands, India. *Proceedings of the International Academy of Ecology and Environmental Sciences* **3**, 36–41.
- Sreeraj CR and Raghunathan C** (2015) A report on the coral reef dwelling polyclads of Nicobar Islands, India. *Proceedings of the International Academy of Ecology and Environmental Sciences* **5**, 83–88.
- Sreeraj CR, Raghunathan C, Raghuraman R, Dixit S and Venkataraman K** (2015) *Polyclads of Andaman and Nicobar Islands*. Kolakata: Zoological Survey of India, 100 pp.
- Tyler S, Artois T, Schilling S, Hooge M and Bush LF** (eds) (2006–2020) World List of turbellarian worms: Acoelomorpha, Catenulida, Rhabditophora. *Pseudoceros* Lang, 1884. Accessed through: World Register of Marine Species. <http://www.marinespecies.org/aphia.php?p=tax-details&id=142241> (Accessed 15 August 2020).
- Velasquez X, Bolaños DM and Benayahu Y** (2018) New records of cotylean flatworms (Platyhelminthes: Polycladida Rhabditophora) from coastal habitats of Israel. *Zootaxa* **4438**, 237–260.
- Verrill AE** (1901) Additions to the fauna of the Bermudas from the Yale Expedition of 1901, with notes on other species. *Transactions of the Connecticut Academy of Arts and Sciences* **11**, 15–62.
- Yeri M and Kaburaki T** (1920) Notes on two new species of Japanese polyclads. *Annotationes Zoologicae Japonenses* **9**, 591–598.